



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

the time, but shortly afterwards several more were found in the hardwood trees on the brow of a hill in the immediate vicinity. In both instances the birds, although near water, were on comparatively high ground and at some distance from the swampy coverts which we would have expected them to affect.—H. F. MOORE, *Philadelphia, Pa.*

Dendroica tigrina at Iowa City, in November.—A small bird taken at Iowa City, Iowa, November 27, proved to be a Cape May Warbler (*Dendroica tigrina*). It was feeding in the top of a pine tree in one of the door-yards at that place, where it was shot. It was in immature plumage, which was very much soiled with pine resin; but otherwise was in good condition.—CHARLES R. KEYES, *Des Moines, Iowa.*

Bird Notes from Toronto.—A male *Sturnella magna* was collected Feb. 21, 1881, by Mr. Jas. Helliwell, at Highland Creek, about fifteen miles east of Toronto. The bird was in fine plumage and in fairly good condition. He had his 'home' in a dense thicket in a deep ravine, through which ran a 'Spring Creek' (which did not freeze during the winter), about a mile from a barn-yard which he visited almost daily, feeding on sweepings and pickings from manure. The bird was carefully dissected but no wound or injury of any kind could be found. The gizzard contained a few small pieces of gravel, a few grains of oats, and pickings from cow dung. This is believed to be the first record of this species wintering north of Lake Ontario.

A male *Melospiza fasciata* was collected Feb. 2, 1886, by Mr. Wm. Squires, while feeding on amaranthus seeds in a garden in St. Matthews-ward, Toronto. Snow ten inches in depth. Another specimen was taken Jan. 31, 1887, by Mr. Daniel G. Cox, in a ravine in St. James Cemetery, Toronto, in a willow thicket densely grown with goldenrod (*Solidago*). Snow six inches deep.

April 4, 1886. *Merula migratoria* Linn. Gizzard contained three hipps of *Rosa blanda* and one larva of *Pyrrharctia isabella*. Ground frozen. It is not usual for any bird to feed on the larvæ of this moth.

A male *Icterus spurius* was collected May 13, 1887, while pursuing insects through willow blossoms, just east of the city limits. Believed to be the first authentic record of the occurrence of this bird at Toronto.

A male European Goldfinch (*Carduelis elegans*) was collected May 21, 1887, by Mr. Daniel S. Cox, about a mile north of the city limits—one out of four—while resting on the top of a beech tree. The remaining three flew off in a northerly direction. The birds were evidently in a natural condition and migrants from the south, doubtless from the New York colony.—WILLIAM BRODIE, *Toronto, Can.*